

Playing with Triangular Numbers

Enrique Treviño



LAKE FOREST
COLLEGE

Chico State Colloquium
December 12, 2018

Dipika Subramaniam




Paul Pollack



Triangular Numbers

What are triangular numbers?



Triangular Numbers

The n -th triangular number, Δ_n is $\frac{n(n+1)}{2}$

Classic proof:

$$\Delta_n = 1 + 2 + \cdots + n$$

$$\Delta_n = n + (n-1) + \cdots + 1$$

$$2\Delta_n = (n+1) + (n+1) + \cdots + (n+1)$$

$$2\Delta_n = n(n+1)$$

$$\Delta_n = \frac{n(n+1)}{2}.$$

Combinatorial Proof:

$$\sum_{i=1}^n i = \sum_{i=1}^n \sum_{j=0}^{i-1} 1 = \binom{n+1}{2}.$$

Probabilistic Proof

Let X be the sum of two uniformly-distributed n -sided dice.

$$\mathbb{P}[X = k] = \begin{cases} \frac{(k-1)}{n^2}, & 2 \leq k \leq n+1 \\ \frac{n-i+1}{n^2}, & k = n+i \text{ with } 2 \leq i \leq n \end{cases}$$

Since $2 \leq X \leq 2n$, then

$$1 = \sum_{k=2}^{2n} \mathbb{P}[X = k] = \sum_{k=2}^{n+1} \frac{k-1}{n^2} + \sum_{i=2}^n \frac{n-i+1}{n^2}$$

$$1 = \left(\frac{1}{n^2} + \frac{2}{n^2} + \cdots + \frac{n}{n^2} \right) + \left(\frac{n-1}{n^2} + \frac{n-2}{n^2} + \cdots + \frac{1}{n^2} \right)$$

$$n^2 = (1 + 2 + \cdots + n) + (1 + 2 + \cdots + (n-1))$$

$$n^2 = 2(1 + 2 + \cdots + n) - n.$$

Playing with Triangular Numbers


$$1 + 3 + 6 = 10$$

McMullen, inspired by this, asked himself:

- For which k can we find k consecutive triangular numbers that add up to be a triangular number?
- Can we find the solutions?

McMullen showed there are infinitely many solutions for $k = 2, 3, 5$, but no solutions for $k = 4$.

Example



$k = 4$ case

Elementary manipulations show that the sum of the k -consecutive triangular numbers starting at Δ_n is Δ_m whenever

$$(2m+1)^2 - k(2n+k)^2 = \frac{(k-1)(k^2+k-3)}{3}.$$

When $k = 4$ we get

$$(2m+1-4n-8)(2m+1+4n+8) = 17.$$

From which

$$(m, n) = (4, 0), (4, -4), (-5, 0), \text{ and } (-5, -4).$$

Theorem

Let $k > 4$ be a square. Then there exist k consecutive triangular numbers that add up to make a bigger triangular number.

Useful Factorization

Let $k = a^2$.

Then

$$(2m+1)^2 - a^2(2n+a^2)^2 = \frac{a^6 - 4a^2 + 3}{3}.$$

$$(2m+1+2na+a^3)(2m+1+2na+a^3) = \frac{(a+1)(a-1)(a^4+a^2-3)}{3}$$

k an even square

$$(2m + 1 - 2na - a^3)(2m + 1 + 2na + a^3) = \frac{a^6 - 4a^2 + 3}{3}$$

Solving

$$2m + 1 - 2na - a^3 = 1$$

$$2m + 1 + 2na + a^3 = \frac{a^6 - 4a^2 + 3}{3}$$

yields

$$m = \frac{(a^4 - 4)(a^2)}{12}$$
$$n = \frac{a(a^4 - 6a - 4)}{12}$$

k an odd square

We'll consider three cases:

$$a \equiv 0 \pmod{3}$$

$$a \equiv 1 \pmod{3}$$

$$a \equiv 2 \pmod{3}$$

$$a \equiv 1 \pmod{3} \text{ or } a \equiv 0 \pmod{3}$$

$$(2m + 1 + 2na + a^3)(2m + 1 + 2na + a^3) = \frac{(a + 1)(a - 1)(a^4 + a^2 - 3)}{3}$$

Solving

$$2m + 1 - 2na - a^3 = a + 1$$

$$2m + 1 + 2na + a^3 = \frac{(a - 1)(a^4 + a^2 - 3)}{3}$$

yields

$$m = \frac{a^2(a - 1)(a^2 + 1)}{12}$$
$$n = \frac{(a + 2)(a - 3)(a^2 + 1)}{12}$$

$$a \equiv 2 \pmod{3} \text{ or } a \equiv 0 \pmod{3}$$

$$(2m + 1 + 2na + a^3)(2m + 1 + 2na + a^3) = \frac{(a + 1)(a - 1)(a^4 + a^2 - 3)}{3}$$

Solving

$$2m + 1 - 2na - a^3 = a - 1$$

$$2m + 1 + 2na + a^3 = \frac{(a + 1)(a^4 + a^2 - 3)}{3}$$

yields

$$m = \frac{a^5 + a^4 + a^3 + a^2 - 12}{12}$$

$$n = \frac{(a + 3)(a - 2)(a^2 + 1)}{12}$$

$$k = 6$$

Recall

$$(2m+1)^2 - k(2n+k)^2 = \frac{(k-1)(k^2+k-3)}{3}.$$

When $k = 6$:

$$x^2 - 6y^2 = 65.$$

Therefore $x^2 \equiv 6y^2 \pmod{13}$. But

$$\left(\frac{6}{13}\right) = -1.$$

Therefore, there are no solutions for $k \equiv 6 \pmod{13}$.

A sufficient condition for k

Lemma

Let $q > 3$ be a prime number. Suppose that $k \in \mathbb{Z}$ is such that

- ① *k is not a square modulo q ,*
- ② *$q \parallel k^2 + k - 3$.*

Then there are no k consecutive triangular numbers that add up to a triangular number.

Example:

If $k \equiv 45 \pmod{53}$ and $k \not\equiv 2430 \pmod{53^2}$. There are 52 residues modulo 53^2 which satisfy these conditions.

Main Theorem

Theorem

Let $K(x)$ be the number of k 's less than x that have solutions. Then:

$$\sqrt{x} \leq K(x) \ll \frac{x}{\sqrt{\log(x)}}.$$

Finding q such that $q \parallel k^2 + k - 3$ and $(k/q) = -1$

If $q \neq 13$, $k^2 + k - 3 \equiv 0 \pmod{q}$ has two distinct solutions k_1, k_2 whenever $(13/q) = 1$. We then have three possibilities

- 1 Both k_1, k_2 are squares modulo q .
- 2 One of k_1, k_2 is a square and the other one isn't.
- 3 Neither k_1, k_2 are squares modulo q .

A pair of important sets of primes

Let \mathcal{A} be the set of primes q for which we have k_1, k_2 both nonsquares modulo q .

Let \mathcal{B} be the set of primes q for which exactly one of k_1, k_2 is a square modulo q .

If $q \in \mathcal{A}$, then the proportion of residues modulo q^2 one must avoid are

$$2\frac{q-1}{q^2} = \frac{2}{q} - \frac{2}{q^2}.$$

If $q \in \mathcal{B}$, then the proportion of residues modulo q^2 one must avoid are

$$\frac{q-1}{q^2} = \frac{1}{q} - \frac{1}{q^2}.$$

Quantifying the proportion of primes in \mathcal{A} , \mathcal{B}

Consider $f(x) = x^4 + x^2 - 3$. Let's analyze how $f(x)$ might factor in \mathbb{Z}_q . There are several possibilities

- $(1,1,1,1)$
- $(1,1,2)$
- $(2,2)$
- 4

Primes in \mathcal{B} would split as $(1,1,2)$.

Primes in \mathcal{A} would be primes that are squares modulo 13 and that don't split as $(1,1,2)$ or $(1,1,1,1)$.

Chebotarev Density Theorem

Theorem

Suppose that $f(x) \in \mathbb{Z}[x]$ is monic and irreducible over \mathbb{Q} , with $\deg f(x) = n$. Let \mathbb{L} be the splitting field of $f(x)$ over \mathbb{Q} . Fix a partition $\langle k_1, \dots, k_r \rangle$ of n (that is, a tuple of positive integers $k_1 \geq k_2 \geq \dots \geq k_r$ with $k_1 + \dots + k_r = n$). Let δ be the proportion of elements of $\text{Gal}(\mathbb{L}/\mathbb{Q})$ which, when viewed as permutations on the roots of $f(x)$, have cycle type $\langle k_1, \dots, k_r \rangle$. For all but finitely many primes p , the polynomial $f(x)$ factors as a product of distinct monic irreducible polynomials modulo p , and δ is the proportion of primes for which these irreducibles have degrees k_1, \dots, k_r .

Chebotarev in our problem

Consider $f(x) = x^4 + x^2 - 3$. f is irreducible over \mathbb{Q} , let \mathbb{L} be the splitting field of f over \mathbb{Q} , then $\text{Gal}(\mathbb{L}/\mathbb{Q})$ is isomorphic to

$$\{(1), (1324), (12)(34), (1423), (34), (13)(24), (12), (14)(23)\}.$$

- 1 of the 8 elements decompose as $(1,1,1,1)$
- 3 of the 8 elements decompose as $(2,2)$
- 2 of the 8 elements decompose as $(1,1,2)$
- 2 of the 8 elements decompose as (4)

The proportion of primes $q \in \mathcal{B}$ is $2/8 = 1/4$.

The proportion of primes $q \in \mathcal{A}$ is $1/2 - 1/8 - 2/8 = 1/8$.

There are several residues modulo certain squares of primes that must be avoided for k to be able to yield solutions.

We then get the following upper bound heuristic:

$$\begin{aligned} K(x) &\ll x \prod_{\substack{q \leq x \\ q \in \mathcal{A}}} \left(1 - \frac{2}{q} + \frac{2}{q^2}\right) \prod_{\substack{q \leq x \\ q \in \mathcal{B}}} \left(1 - \frac{1}{q} + \frac{1}{q^2}\right) \\ &\ll x \left(\frac{1}{(\log^{1/8} x)^2} \right) \left(\frac{1}{\log^{1/4}(x)} \right) \\ &\ll \frac{x}{\sqrt{\log x}}. \end{aligned}$$

Non-Chebotarev (weaker) proof

One could avoid the use of Chebotarev to get that $K(x) = o(x)$.
Namely, the primes in \mathcal{B} can be characterized as primes q satisfying

$$\left(\frac{13}{q}\right) = 1 \quad \& \quad \left(\frac{-3}{p}\right) = -1.$$

The primes in \mathcal{B} have proportion $1/4$.

Then

$$K(x) \ll \frac{x}{(\log(x))^{1/4}}.$$

Main Theorem

Theorem

Let $K(x)$ be the number of k 's less than x that have solutions. Then:

$$\sqrt{x} \leq K(x) \ll \frac{x}{\sqrt{\log(x)}}.$$

On the hunt for a lower bound

Our goal is solving

$$(2m+1)^2 - k(2n+k)^2 = \frac{(k-1)(k^2+k-3)}{3}.$$

Let p be a prime number satisfying:

- 1 $p \equiv 7 \pmod{24}$
- 2 $p^2 + p - 3$ is not divisible by any prime q for which $p \pmod{q}$ is a nonsquare
- 3 $\mathbb{Q}(\sqrt{p})$ has class number 1.

Then there exist p consecutive triangular numbers that add up to a triangular number.

$$k = 127$$

We want to solve

$$(2m + 1)^2 - 127(2n + 127)^2 = 682626 = 2 \times 3 \times 7 \times 16253.$$

- $127 \equiv 1 \pmod{42}$
- $541^2 \equiv 127 \pmod{16253}$
- $\mathbb{Q}(\sqrt{127})$ has class number 1.

Let $q \in \{2, 3, 7, 16253\}$. There exists $x_q + y_q\sqrt{127}$ with norm q .

- $x_2 = 2175, y_2 = 193$
- $x_3 = 293, y_3 = 26$
- $x_7 = 45, y_7 = 4$
- $x_{16253} = 2325, y_{16253} = 206$

Solution to $k = 7$

$$(45 + 4\sqrt{127})(293 + 26\sqrt{127})(2175 + 193\sqrt{127})(2325 + 206\sqrt{127}) \\ = 533462754763 + 47337164797\sqrt{127}$$

Let

$$x = 533462754763, \quad y = 47337164797.$$

Then

$$x^2 - 127y^2 = 682626.$$

We want to solve $2m + 1 = x$ and $2n + 127 = y$.

$$m = 266731377381, \quad n = 23668582335.$$

$$\Delta_n + \Delta_{n+1} + \cdots + \Delta_{n+126} = \Delta_m.$$

Conjecture

Let \mathcal{P} be the set of prime numbers p satisfying

- ① $p \equiv 7 \pmod{24}$
- ② $p^2 + p - 3$ is not divisible by any prime q for which $p \pmod{q}$ is a nonsquare
- ③ $\mathbb{Q}(\sqrt{p})$ has class number 1.

The proportion of such primes is 75.45%.

This suggests

$$K(x) \gg \frac{x}{\log^{3/2}(x)}.$$

Thank you!